

Prevalence of Sexually Transmitted Infections and Predictors for Loss to Follow Up among Marginalized Homeless and Migrant Communities: a Cross-Sectional Study



ORIGINAL RESEARCH

Ju[ubiquity press

FRANCESCO VLADIMIRO SEGALA

ROBERTA NOVARA

GIANFRANCO PANICO

RENATO LAFORGIA

LUCIA RAHO

MARCELLA SCHIAVONE

GIOVANNI CIVILE

NICOLE LAFORGIA

STEFANO DI GREGORIO

GIACOMO GUIDO

MARIANGELA CORMIO

ANGELO DARGENIO

ROBERTA PAPAGNI

ANGELO L'ERARIO

LUCA L'ERARIO

VALENTINA TOTARO

VITO SPADA

LAURIANA VALENTINI

LUISA FRALLONARDO

ROSSANA LATTANZIO

CARMINE FALANGA

GIOVANNI PUTOTO

ANNALISA SARACINO

FRANCESCO DI GENNARO

CORRESPONDING AUTHOR:

Roberta Novara, MD

Clinic of Infectious Diseases,
Department of Precision and
Regenerative Medicine and
Ionian Area – (DiMePRE-J),
University of Bari 'Aldo Moro,'
Piazza Giulio Cesare n. 11 Cap,
Bari, Italy

robertanovara@gmail.com

*Author affiliations can be found in the back matter of this article

KEYWORDS:

STI; homeless; migrants;
screening; return rate;
healthcare

ABSTRACT

Background: In Europe and Italy, marginalized communities have a higher risk for both contracting sexually transmitted infections (STI) and progressing towards adverse outcomes.

Objectives: This study focuses on the screening of HIV, HBV, HCV, and syphilis among homeless individuals and agricultural migrant workers living in Apulia, Italy. It aims to assess STI prevalence and investigate factors that might hinder return to collect test results. In addition, it explores STI knowledge, attitudes, and practices among these vulnerable populations.

Methods: A cross-sectional study was conducted from September 1, 2022, to September 30, 2023. Participants were recruited from community health centers and migrant camps. Blood tests for HBV, HCV, HIV, and syphilis were performed, and Knowledge, Attitude, and Practices (KAP) survey were conducted via face-to-face interviews. Descriptive and logistic regression analyses were used to assess factors influencing the return for test results.

TO CITE THIS ARTICLE:

Segala FV, Novara R, Panico G, Laforgia R, Raho L, Schiavone M, Civile G, Laforgia N, Di Gregorio S, Guido G, Cormio M, Dargenio A, Papagni R, L'Erario A, L'Erario L, Totaro V, Spada V, Valentini L, Frallonardo L, Lattanzio R, Falanga C, Putoto G, Saracino A, Di Gennaro F. Prevalence of Sexually Transmitted Infections and Predictors for Loss to Follow Up among Marginalized Homeless and Migrant Communities: a Cross-Sectional Study. *Annals of Global Health*. 2024; 90(1): 25, 1–13. DOI: <https://doi.org/10.5334/aogh.4388>

organization Doctors with Africa CUAMM (University College for Aspiring Medical Missionaries). This organization has been actively engaged in assisting the agricultural migrant worker communities since 2015. STI and KAP screening surveys were delivered concurrently by face-to-face interviews.

Study sites included four community health centers for homeless people situated in the City of Bari and five resettlement camps located in the municipality of Foggia, so-called ‘ghettos,’ where agricultural migrant workers (AMW) reside. A more detailed description of sites included in the study is provided in the supplementary file.

The eligible population included all adult individuals who were beneficiaries or residents in the mentioned facilities throughout the study period. No exclusion criteria were applied for this study. Participants had the option to withdraw from the study at any time, and their decision did not impact their medical care.

Blood test result delivery was scheduled approximately 10 days after screening, and the failure to return for results rate (FTR) was assessed. Participants who tested positive for any of the infections included in the study were referred for further diagnostic evaluation and treatment to the Clinic of Infectious Diseases of the Polyclinic of Bari. Positive-screened participants received the same standard of care, in accordance with national and international guidelines [10–13].

QUESTIONNAIRE DEVELOPMENT AND DATA COLLECTION

Questionnaire development was informed by literature review and administered through face-to-face interviews conducted by nurses and trained doctors. Every interaction with the subjects involved in the study was facilitated by linguistic and cultural mediators, when needed. To minimize the social desirability bias, interviewers provided clear instructions at the beginning of the questionnaire, explaining the purpose of the survey, the importance of honest responses, and reassuring participants about data confidentiality.

The questionnaire was organized into the following sub-sections:

- Socio-demographic information (age, education, occupation, marital status, how long they were homeless, possible employment contract, residence document in Italy): 6 questions.
- Risk factors for STI: multiple sexual partners, use of condoms, smoking and alcohol habits, chem-sex (defined as “voluntary intake of psychoactive drugs with the intention of facilitating and/or enhancing sexual activity [14]”) and medical history (including symptoms of TB and other communicable-non-communicable diseases): 6 questions.
- Knowledge, attitude, and practices on HIV, HBV, HCV, and Syphilis: 25 questions.

All study participants underwent the following blood tests:

- Serologies for HBV (HBsAg, anti-HBs, and anti-HBc), HCV (anti-HCV), human immunodeficiency virus (HIV), and syphilis (VDRL and TPHA).
- The study participants, resulting positive for the screening, performed confirmation tests according to the standard of care: HBV-DNA for HBV, HCV-RNA and genotype for HCV, HIV-RNA, immune cell count, and resistance profile for HIV.

All data were entered into a secured online platform (Kobo Ttoolbox [15]), and a quality control check of the data entry was performed before data analysis.

STATISTICAL ANALYSIS

A descriptive analysis was performed to define the distribution of the characteristics of the sample. Continuous data were summarized as median and interquartile range (IQR). Return For Test Results (RTR) was identified as the dependent variable. World regions were classified according to the United Nation Geoscheme [16]. The distribution of continuous variables was assessed with the Shapiro-Wilk normality test, and the Mann-Whitney U test was used to compare groups for continuous variables. A chi-squared test (with the Fisher’s correction if less than five cases were

present in a cell) was applied for categorical variables. A logistic regression model was implemented as follows. RTR was considered as a dependent variable and each of the available factors at the baseline evaluation as independent variables (univariate analysis). The effect sizes were reported as odds ratio (OR) with 95% confidence interval (CI). All tests were 2-sided, and a p-value <0.05 was considered statistically significant. The model included a set of clinically relevant candidate predictors and variables resulted to be significant at the bivariate analysis. Statistical analyses were performed using R Statistical Software (v4.1.3; R Core Team 2021) in R Studio Version [17].

ETHICAL CONSIDERATIONS

Prior to recruitment, written informed consent (prepared in 4 languages: Italian, French, English, and Arabic) was obtained from all participants. The objectives of the study, along with methods used and data confidentiality, were explained with the aid of cultural mediators, when needed. The study has been approved by Ethics Committee ‘Azienda Ospedaliero-Universitaria Consorziale Policlinico,’ protocol number DG1563. During the screening process, we actively worked to establish and enhance personalized pathways for connecting individuals to healthcare through various social and healthcare services. We raised awareness about the potential progression of HIV and liver diseases, as well as the risk of transmission to others, by directing these individuals to the Infectious Disease Clinic at the Policlinic of Bari for counseling, diagnosis, and appropriate treatment.

RESULTS

CHARACTERISTICS OF THE POPULATION

From September 1, 2022, to September 30, 2023, a total of 149 participants were recruited in the study, of which 64 (42.9%) were agricultural migrant workers and 85 (57.1%) were homeless, with a median age of 35 (IQR: 27–45) years. Characteristics of the sample stratified for study subpopulation and RTR are displayed in Table 1. Males accounted for 87.5% (n = 56) of agricultural migrant workers and 75.3% (n = 64) of homeless people. The most frequently reported comorbidities were diabetes (8.1%, n = 12), hypertension (6%, n = 9), and respiratory pathologies (4%, n = 6). Among agricultural migrant workers, countries of origin were located mainly in sub-Saharan Africa (64.1%, n = 41) and North Africa (20.3%, n = 13), while 32.9% (n = 28) and 31.8% (n = 27) of homeless people were originally from Sub-Saharan Africa and Italy. Educational level was low, with 64.4% (n = 96) of the included subjects reporting only primary school or no formal education. Overall, 69.1% (n = 103) of the participants did not have access to primary healthcare, and this rate was alarmingly high among people residing in the Apulian ghettos (85.9% n = 55 vs 56.5% n = 48, p < 0.001). Apart from country of origin (p < 0.001) and access to care, the migrant and homeless subpopulation differed significantly on the following variables: reporting chem-sex (p = 0.049) and HCV and HIV knowledge (p < 0.001 and p = 0.02)—with homeless people scoring better.

Table 1 Characteristics of the population stratified for subpopulation and return rate.

	OVERALL (N = 149)	SUB-POPULATION			RETURN FOR TEST RESULTS		
		AGRICULTURAL MIGRANT WORKERS (N = 64)	HOMELESS PEOPLE (N = 85)	P-VALUE FOR SETTING	NOT RETURNED (N = 74)	RETURNED (N = 75)	P-VALUE FOR RETURNING
Age, median [Q1, Q3]	35 [27, 45]	34 [26, 42]	35 [29, 48]	0.328	33.0 [26.3, 41.8]	36.0 [28.5, 50.0]	0.171
Gender							
F	29 (19.5%)	8 (12.5%)	21 (24.7%)	0.0982	11 (14.9%)	18 (24.0%)	0.23
M	120 (80.5%)	56 (87.5%)	64 (75.3%)		63 (85.1%)	57 (76.0%)	
Sub-populations							
Agricultural migrant workers	64 (43.0%)	–	–	–	38 (51.4%)	26 (34.7%)	0.0585
Homeless people	85 (57.0%)	–	–		36 (48.6%)	49 (65.3%)	

(Contd.)

	OVERALL (N = 149)	SUB-POPULATION			RETURN FOR TEST RESULTS		
		AGRICULTURAL MIGRANT WORKERS (N = 64)	HOMELESS PEOPLE (N = 85)	P-VALUE FOR SETTING	NOT RETURNED (N = 74)	RETURNED (N = 75)	P-VALUE FOR RETURNING
Marital Status							
Divorced	11 (7.4%)	3 (4.7%)	8 (9.4%)	0.442	3 (4.1%)	8 (10.7%)	0.478
In an open relationship	4 (2.7%)	2 (3.1%)	2 (2.4%)		2 (2.7%)	2 (2.7%)	
Single	84 (56.4%)	34 (53.1%)	50 (58.8%)		44 (59.5%)	40 (53.3%)	
Married	49 (32.9%)	25 (39.1%)	24 (28.2%)		25 (33.8%)	24 (32.0%)	
Missing	1 (0.7%)	0 (0%)	1 (1.2%)		0 (0%)	1 (1.3%)	
Region of origin							
Eastern Europe	10 (6.7%)	6 (9.4%)	4 (4.7%)	<0.001	4 (5.4%)	6 (8.0%)	0.25
Middle East	10 (6.7%)	1 (1.6%)	9 (10.6%)		8 (10.8%)	19 (25.3%)	
North Africa	24 (16.1%)	13 (20.3%)	11 (12.9%)		4 (5.4%)	6 (8.0%)	
Other	2 (1.3%)	1 (1.6%)	1 (1.2%)		14 (18.9%)	10 (13.3%)	
Southeast Asia	7 (4.7%)	2 (3.1%)	5 (5.9%)		1 (1.4%)	1 (1.3%)	
Sub-Saharan Africa	69 (46.3%)	41 (64.1%)	28 (32.9%)		3 (4.1%)	4 (5.3%)	
Italy	27 (18.1%)	0 (0%)	27 (31.8%)		40 (54.1%)	29 (38.7%)	
Education							
Primary school	56 (37.6%)	29 (45.3%)	27 (31.8%)	0.0872	33 (44.6%)	23 (30.7%)	0.373
Lower secondary school	36 (24.2%)	13 (20.3%)	23 (27.1%)		15 (20.3%)	21 (28.0%)	
Higher secondary school	14 (9.4%)	4 (6.3%)	10 (11.8%)		5 (6.8%)	9 (12.0%)	
University	3 (2.0%)	3 (4.7%)	0 (0%)		1 (1.4%)	2 (2.7%)	
No formal education	40 (26.8%)	15 (23.4%)	25 (29.4%)		20 (27.0%)	20 (26.7%)	
Comorbidities							
Diabetes	12 (8.1%)	2 (3.1%)	10 (11.8%)	0.106	4 (5.4%)	8 (10.7%)	0.379
Hypertension	9 (6.0%)	2 (3.1%)	7 (8.2%)	0.343	4 (5.4%)	5 (6.7%)	1
Cardiopathy	4 (2.7%)	0 (0%)	4 (4.7%)	0.212	1 (1.4%)	3 (4.0%)	0.622
Pneumopathy	6 (4.0%)	0 (0%)	6 (7.1%)	0.0803	3 (4.1%)	3 (4.0%)	1
Neuropsychiatric disorder	4 (2.7%)	0 (0%)	4 (4.7%)	0.212	2 (2.7%)	2 (2.7%)	1
Cancer	2 (1.3%)	1 (1.6%)	1 (1.2%)	1	0 (0%)	2 (2.7%)	0.482
TB History	3 (2.0%)	1 (1.6%)	2 (2.4%)	1	1 (1.4%)	2 (2.7%)	1
Chem sex	14 (9.4%)	2 (3.1%)	12 (14.1%)	0.0494	4 (5.4%)	10 (13.3%)	0.177
Missing	1 (0.7%)	1 (1.6%)	0 (0%)		1 (1.4%)	0 (0%)	
Consistent condom use	21 (14.1%)	10 (15.6%)	11 (12.9%)	0.819	9 (12.2%)	12 (16.0%)	0.662
Access to primary healthcare	46 (30.9%)	9 (14.1%)	37 (43.5%)	<0.001	15 (20.3%)	31 (41.3%)	0.009
Fear of community exclusion in case of HIV positivity	88 (59.1%)	37 (57.8%)	51 (60.0%)	0.92	43 (58.1%)	45 (60.0%)	0.946

(Contd.)

	OVERALL (N = 149)	SUB-POPULATION			RETURN FOR TEST RESULTS		
		AGRICULTURAL MIGRANT WORKERS (N = 64)	HOMELESS PEOPLE (N = 85)	P-VALUE FOR SETTING	NOT RETURNED (N = 74)	RETURNED (N = 75)	P-VALUE FOR RETURNING
Fear of community exclusion in case of HCV positivity	38 (25.5%)	16 (25.0%)	22 (25.9%)	1	16 (21.6%)	22 (29.3%)	0.372
Being tested for HIV or HCV in the past	38 (25.5%)	14 (21.9%)	24 (28.2%)	0.489	18 (24.3%)	20 (26.7%)	0.889
HCV Knowledge Score, median [Q1, Q3]	0 [0, 2]	0 [0, 1]	1 [0, 3]	0.001	0 [0, 1]	0 [0, 3]	0.04
HIV Knowledge Score, median [Q1, Q3]	2 [0, 4]	2 [0, 3]	3 [1, 4]	0.019	2 [0, 3]	2 [0, 4]	0.512
Infections							
HIV	3 (2.0%)	2 (3.1%)	1 (1.2%)	0.803	1 (1.4%)	2 (2.7%)	1
Syphilis	5 (3.4%)	2 (3.1%)	3 (3.5%)	1	5 (6.8%)	0 (0%)	0.0665
HCV	14 (9.4%)	6 (9.4%)	8 (9.4%)	1	12 (16.2%)	2 (2.7%)	0.01
HBV Infection	21 (14.1%)	10 (15.6%)	11 (12.9%)	0.819	11 (14.9%)	10 (13.3%)	0.974
Any infection	37 (24.8%)	17 (26.6%)	20 (23.5%)	0.816	24 (32.4%)	13 (17.3%)	0.052
More than one infection	6 (4.0%)	3 (4.7%)	3 (3.5%)	1	5 (6.8%)	1 (1.3%)	0.205
Vaccinated against HBV	21 (14.1%)	5 (7.8%)	16 (18.8%)	0.078	7 (9.5%)	58 (77.3%)	0.138
Missing	3 (2.0%)	0 (0%)	3 (3.5%)		0 (0%)	14 (18.7%)	
Returned for test results	75 (50.3%)	26 (40.6%)	49 (57.6%)	0.058	-	-	-

KNOWLEDGE, ATTITUDES, AND PRACTICES TOWARDS SEXUALLY TRANSMITTED INFECTIONS

Concerning HIV/AIDS, only 60% (n = 90) of the recruited individuals were aware that the disease can be transmitted through sexual contact. Additionally, 45.7% (n = 65) were not aware of the protection offered by condoms. A notable 34.9% (n = 52) agreed or strongly agreed that HIV/AIDS might be transmitted by sharing utensils or having a meal together. Moreover, over 75% (n = 111) of the sample did not agree with the statement ‘thanks to HIV treatment, people can lead a high-quality life and not be contagious anymore.’ Awareness of Hepatitis C was notably low, with limited knowledge about its modes of transmission, potential complications, and available treatment options in both subpopulations. Details about the distribution of responses to the KAP questionnaire are provided in **Supplementary Table 1**. Both HIV (p < 0.001) and HCV (p = 0.02) knowledge were significantly lower among agricultural migrant workers ([Table 1](#)).

Regarding the perception of stigma, the majority (59.1%, n = 88) of participants expressed fear that a positive HIV test would lead to social exclusion, while only 25.5% (n = 38) held similar concerns about a positive HCV test. In terms of practices, the majority of participants (85.9%, n = 128) reported inconsistent condom use, while 14.1% (n = 12) of homeless individuals stated that they had engaged in chem-sex.

SEXUALLY TRANSMITTED INFECTIONS SCREENING

The total number of infections detected is illustrated in [Figure 1](#). Overall, out of 149 participants screened, 3 tested positive for HIV (2%), 14 for HCV (9.4%), 21 (14.1%) for HBV, and 5 (3.4%) for syphilis. Only 14% were vaccinated for HBV. HBV serology results ([Figure 2](#)) indicate that most of the study participants have neither been vaccinated against HBV nor have they ever been exposed

to the virus. Approximately one out of four people (24.8%, n = 37) were positive for at least one infection, while three people out of four (74.5%, n = 111) reported that they had never undergone prior testing for these infections.




Figure 1 Distribution of infections among migrants and homeless people.




Figure 2 HBV serologies stratified per subpopulation.

MONO AND MULTIVARIABLE ANALYSIS FOR RETURN RATE FOR TEST RESULTS

Overall, only half of the screened participants (50.3%, n = 75) returned to collect test results. As shown in [Table 1](#), at univariate analysis, people who had access to primary healthcare were significantly associated with returning (41.3% n = 31 vs 20.3% n = 15, p < 0.009), as well as people who performed better at HCV knowledge questionnaire (p = 0.04). Alarming, people whose results indicated infection with HCV or with at least one STI were more likely to be lost to follow-up.

At multivariable analysis, the only factor independently associated with reduced odds of returning to collect test results was being positive to HCV ([Table 2](#)).

FACTOR	aOR	lowCI	highCI	pVALUE
(Intercept)	0.845	0.398	1.775	0.657
Living in Apulian Ghettos	0.697	0.332	1.462	0.337
Having access to primary healthcare	2.101	0.928	4.881	0.077
Good HCV knowledge	2.208	0.911	5.583	0.083
Having contracted HCV	0.136	0.017	0.703	0.028
Having contracted at least one STD	0.870	0.334	2.249	0.773
Fear that positive HIV test would result in social exclusion	1.214	0.596	2.483	0.592

Table 2 Multivariable logistic regression model for return for test results.

DISCUSSION

In our study, which included 85 homeless people and 64 agricultural migrant workers living in Southern Italy, the cumulative prevalence of STI was high, and only half of the overall population returned to collect test results. Overall knowledge of STI modes of transmission was low and, in line with other studies [18, 19], only 14% of the population reported consistent condom use.

In the present study, almost one homeless person out of ten reported having been engaged in chem-sex. This is concerning, because it is likely a proxy of the high rate of substance use disorder—and possibly also other mental health disorders—described in this and other marginalized populations [20]. Indeed, marginalization is correlated with increased high-risk behaviors [21] as well as higher disease burden and mortality [22]. In our population, the cumulative prevalence of STI was 24% and, interestingly, rates were similar among agricultural migrant workers (26.6%) and homeless people (23.5%), substantially exceeding the prevalence rates observed in the general population [23]. This data might be explained by the fact that the root causes of the disease burden lay in marginalization and social vulnerability themselves, rather than individual misconceptions or diverse cultural background [24], thus underscoring the intricate relationship between socio-economic vulnerability, substance abuse, barriers to healthcare, mental health, and the transmission of these viral infections.

Regarding the perceived stigma of HIV, the majority (59.1%) of the participants were afraid that a positive HIV test would result in social exclusion, and this conviction was likely supported by common misconception about disease route of transmission, as it is highlighted by the high rate of people who believed that HIV/AIDS might be contracted by sharing objects or by having meals together. This data, if combined with high STI prevalence might lead infected people to experience further social isolation, imprisoning them in a vicious circle of disease and marginalization.

Indeed, the main finding of our study was that, alarmingly, half of our population did not come back to collect test results, and the strongest predictor for not returning was having contracted HCV. This phenomenon highlights a critical gap in the healthcare system’s ability to engage and retain marginalized populations in the continuum of care. The situation is significantly exacerbated by the lack of consistent healthcare access, as evidenced by 69.1% of participants not having access to primary healthcare. The implications of this limited access are profound, contributing to delayed diagnosis and treatment, and consequently, to worse health outcomes and continued infection transmission [25].

For policymakers, this situation demands a reevaluation of current health policies and systems. It is crucial to implement strategies that not only increase healthcare accessibility but also build trust and engagement with marginalized communities. Furthermore, there is a demand for integrated care models that address not only the medical needs but also the social determinants of health, such as housing, employment, and mental health services [26]. These models can provide a more holistic approach to healthcare, addressing the root causes of healthcare disparities. Moreover, community-based interventions that involve peers or community leaders could significantly improve engagement and trust in healthcare services. This approach demonstrated to be effective in both migrant [27] and homeless [28] populations.

The present study has several limitations. Primarily, the generalizability of our findings is constrained by the study's specific demographic and geographic focus on homeless individuals and agricultural migrant workers in Southern Italy. Secondly, our study's reliance on self-reported data poses a risk of introducing bias, as participants' responses may be influenced by recall and social desirability biases—including the potential underreporting of high-risk behaviors. Additionally, the cross-sectional design of the study inherently limits our ability to establish causality between the observed factors and the prevalence of STIs. Lastly, the absence of longitudinal follow-up in our study design precludes the assessment of long-term outcomes, which are crucial for understanding the progression of STIs and the effectiveness of interventions over time. These limitations highlight the need for further research with diverse populations and marginalized communities, preferably with longitudinal designs to explore the impact of interventions on health outcomes and retention in care.

CONCLUSIONS

Among marginalized homeless and migrant communities, the prevalence of STIs and the failure to return to collect test results were high, and people who contracted HCV were more likely to be lost to follow-up. There is an urgent need to design interventions and policies that address vulnerable communities and explicitly target marginalization and socio-economic inequalities as key determinants of population health.

ADDITIONAL FILE

The additional file for this article can be found as follows:

- **Supplementary file.** Description of study sites and Supplementary Table 1. DOI: <https://doi.org/10.5334/aogh.4388.s1>

ACKNOWLEDGEMENTS

The study has been realized by the collaboration and the support of different non-governmental organization and institutions:

- Doctors with Africa CUAMM, an Italian non-governmental organization that has been present in the ghettos of Foggia since 2015, guaranteeing primary healthcare and awareness-raising activities. For the purposes of the project, it guaranteed logistical support and volunteer medical staff;
- Municipality of Bari, department of Welfare, personal services, social inclusion and fight against poverty, reception, and integration. It is the key department in social services dedicated to the homeless such as dormitories, day services, and canteens. He gave patronage to the Project;
- The School of Specialization in Infectious and Tropical Diseases of the University of Bari, which was the infectious disease and epidemiological partner of our screening activity. For the purposes of the project, the school guaranteed the human resources with the infectious disease skills used in the administration of the tests and the questionnaire;

4. **Ralli M, Cedola C, Urbano S, Morrone A, Ercoli L.** Homeless persons and migrants in precarious housing conditions and COVID-19 pandemic: peculiarities and prevention strategies. *Eur Rev Med Pharmacol Sci.* 2020; 24(18): 9765–9767. DOI: https://doi.org/10.26355/eurev_202009_23071
5. **World Health Organization.** HIV/AIDS in the WHO European Region. Published 2021. Accessed December 1, 2023. <https://www.who.int/europe/news-room/fact-sheets/item/hiv-aids-in-the-who-european-region>.
6. **European Centre for Disease Prevention and Control, World Health Organization.** *HIV/AIDS Surveillance in Europe 2022: 2021 Data.* European Centre for Disease Prevention and Control; 2022. Accessed November 5, 2023. <https://data.europa.eu/doi/10.2900/818446>.
7. **European Centre for Disease Prevention and Control.** Hepatitis B and C testing in the EU/EEA: progress in reaching the elimination targets. Published online 2021. Accessed November 6, 2023. <https://www.ecdc.europa.eu/en/publications-data/hepatitis-b-and-c-testing-eueea-progress-reaching-elimination-targets>.
8. **United Nations.** THE 17 GOALS|Sustainable Development. Published online 2023. Accessed December 2, 2023. <https://sdgs.un.org/goals>.
9. **Mgbako O, Conard R, Mellins CA, Dacus JD, Remien RH.** A systematic review of factors critical for HIV health literacy, ART adherence and retention in care in the U.S. for racial and ethnic minorities. *AIDS Behav.* 2022; 26(11): 3480–3493. DOI: <https://doi.org/10.1007/s10461-022-03680-y>
10. **Consolidated guidelines on HIV prevention, testing, treatment, service delivery and monitoring: recommendations for a public health approach.** Geneva: World Health Organization; 2021. Licence: CC BY-NC-SA 3.0 IGO.
11. **Lampertico P, Agarwal K, Berg T, et al.** EASL 2017 Clinical practice guidelines on the management of hepatitis B virus infection. *J Hepatol.* 2017; 67(2): 370–398. DOI: <https://doi.org/10.1016/j.jhep.2017.03.021>
12. **Pawlotsky JM, Negro F, Aghemo A, et al.** EASL recommendations on treatment of hepatitis C: Final update of the series☆. *J Hepatol.* 2020; 73(5): 1170–1218. DOI: <https://doi.org/10.1016/j.jhep.2020.08.018>
13. **Centers for Disease Control and Prevention.** Sexually Transmitted Infections Treatment Guidelines, 2021. Accessed November 2, 2023. <https://www.cdc.gov/std/treatment-guidelines/default.htm>.
14. **Giorgetti R, Tagliabracchi A, Schifano F, Zaami S, Marinelli E, Busardò FP.** When “chems” meet sex: a rising phenomenon called “chemsex.” *Curr Neuropharmacol.* 2017; 15(5): 762–770. DOI: <https://doi.org/10.2174/1570159X15666161117151148>
15. **KoboToolbox.** KoboToolbox. Accessed March 20, 2023. <https://www.kobotoolbox.org/>.
16. **United Nation Geoscheme.** M49 Standard Version, Geneva. Published 2023. Accessed November 5, 2023. <https://unstats.un.org/unsd/methodology/m49/>.
17. **R core team RF for SC.** A language and environment for statistical computing. Published 2023. Accessed November 5, 2023. <https://www.r-project.org/>.
18. **Totaro V, Patti G, Segala FV, et al.** HIV-HCV Incidence in low-wage agricultural migrant workers sectional study. *Viruses.* Published online 2023; 1–12. DOI: <https://doi.org/10.3390/v15010249>
19. **Di Gennaro F, Lattanzio R, Falanga C, et al.** Low-wage agricultural migrant workers in Apulian Ghettos, Italy: general health conditions assessment and HIV screening. *Trop Med Infect Dis.* 2021; 6(4): 184. DOI: <https://doi.org/10.3390/tropicalmed6040184>
20. **Aldridge RW, Story A, Hwang SW, et al.** Morbidity and mortality in homeless individuals, prisoners, sex workers, and individuals with substance use disorders in high-income countries: a systematic review and meta-analysis. *Lancet Lond Engl.* 2018; 391(10117): 241–250. DOI: [https://doi.org/10.1016/S0140-6736\(17\)31869-X](https://doi.org/10.1016/S0140-6736(17)31869-X)
21. **Montague BT, Vuylsteke B, Buvé A.** Sustainability of programs to reach high risk and marginalized populations living with HIV in resource limited settings: implications for HIV treatment and prevention. *BMC Public Health.* 2011; 11: 701. DOI: <https://doi.org/10.1186/1471-2458-11-701>
22. **Williams SP, Bryant KL.** Sexually transmitted infection prevalence among homeless adults in the United States: a systematic literature review. *Sex Transm Dis.* 2018; 45(7): 494–504. DOI: <https://doi.org/10.1097/OLQ.0000000000000780>
23. **EpiCentro – istituto Superiore di Sanità.** Infezioni sessualmente trasmesse – Aspetti epidemiologici in Italia. Accessed December 21, 2023. <https://www.epicentro.iss.it/ist/epidemiologia-italia>.
24. **Segala FV, Patti G, Olal L, et al.** COVID perceptions among pregnant women living in a Malaria hyperendemic rural region in Uganda: a cross-sectional study. *Am J Trop Med Hyg.* Published online November 6, 2023: tpm230464. DOI: <https://doi.org/10.4269/ajtmh.23-0464>
25. **Riley WJ.** Health disparities: Gaps in access, quality and affordability of medical care. *Trans Am Clin Climatol Assoc.* 2012; 123: 167–174.

26. **Baah FO, Teitelman AM, Riegel B.** Marginalization: Conceptualizing patient vulnerabilities in the framework of social determinants of health – An integrative review. *Nurs Inq.* 2019; 26(1): e12268. DOI: <https://doi.org/10.1111/nin.12268>
27. **Riza E, Kalkman S, Coritsidis A,** et al. Community-based healthcare for migrants and refugees: a scoping literature review of best practices. *Healthcare.* 2020; 8(2): 115. DOI: <https://doi.org/10.3390/healthcare8020115>
28. **Hwang SW, Burns T.** Health interventions for people who are homeless. *Lancet Lond Engl.* 2014; 384(9953): 1541–1547. DOI: [https://doi.org/10.1016/S0140-6736\(14\)61133-8](https://doi.org/10.1016/S0140-6736(14)61133-8)

Segala et al.
Annals of Global Health
DOI: 10.5334/aogh.4388

13

TO CITE THIS ARTICLE:

Segala FV, Novara R, Panico G, Laforgia R, Raho L, Schiavone M, Civile G, Laforgia N, Di Gregorio S, Guido G, Cormio M, Dargenio A, Papagni R, L'Erario A, L'erario L, Totaro V, Spada V, Valentini L, Frallonardo L, Lattanzio R, Falanga C, Putoto G, Saracino A, Di Gennaro F. Prevalence of Sexually Transmitted Infections and Predictors for Loss to Follow Up among Marginalized Homeless and Migrant Communities: a Cross-Sectional Study. *Annals of Global Health.* 2024; 90(1): 25, 1–13. DOI: <https://doi.org/10.5334/aogh.4388>

Submitted: 06 January 2024

Accepted: 08 March 2024

Published: 01 April 2024

COPYRIGHT:

© 2024 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

Annals of Global Health is a peer-reviewed open access journal published by Ubiquity Press.

